

LARGE ROTATOR CUFF REPAIR PROTOCOL

Includes Treatments for:

Open Repair Large Rotator Cuff Tear
Arthroscopic Repair Large Rotator Cuff Tear

Postop:

0-2 Weeks

Sling/swathe at all times, except for exercises

PROM limits to: 90° pure abduction, 20° extension, 70° internal rotation (not behind back)

1. Pendulum exercises 3x/day minimum
2. PROM within pain tolerance
3. Elbow and wrist AROM 4x/day minimum
4. Cryocuff/ ice: Days 1-2 as much as possible, then post activity for pain

2-6 Weeks

Sling/swathe at all times, except for exercises, bathing, or sitting quietly

ROM limits continued

1. PROM within limits and pain tolerance
2. Manual resistance for scapular motions
3. Postural awareness education
4. Gentle soft tissue mobilization and joint mobilization
5. Modalities as indicated for pain or inflammation

6-12 Weeks

Sling for comfort, but may be discontinued

No ROM limits

1. Initiate AAROM (wand and pulleys), unless specified by physician to progress more slowly
2. Joint mobilization for scapular and glenohumeral mobility
3. Gentle exercises for scapular stabilizers and rotator cuff
May include sub-max isometrics, PNF, and supervised UBE
4. Progress AROM with education for quality of motion to avoid shoulder hiking

***Patient must be able to elevate without hiking shoulder to progress**

12-18 Weeks

RESISTIVE EXERCISES HELD AS PER SURGEON INSTRUCTIONS