

BRUNSWICK 207.721.8411 PORTLAND 207.828.2121 SACO 207.710.5504 WINDHAM 207.553.7246

LARGE ROTATOR CUFF REPAIR PROTOCOL

Includes Treatments for:

Open Repair Large Rotator Cuff Tear Arthroscopic Repair Large Rotator Cuff Tear

Postop:

0-2 Weeks Sling/swathe at all times, except for exercises

PROM limits to: 90° pure abduction, 20° extension, 70° internal rotation (not behind back)

- 1. Pendulum exercises 3x/day minimum
- 2. PROM within pain tolerance
- 3. Elbow and wrist AROM 4x/day minimum
- 4. Cryocuff/ ice: Days 1-2 as much as possible, then post activity for pain

2-6 Weeks Sling/swathe at all times, except for exercises, bathing, or sitting quietly ROM limits continued

- 1. PROM within limits and pain tolerance
- 2. Manual resistance for scapular motions
- 3. Postural awareness education
- 4. Gentle soft tissue mobilization and joint mobilization
- 5. Modalities as indicated for pain or inflammation

6-12 Weeks Sling for comfort, but may be discontinued No ROM limits

- 1. Initiate AAROM (wand and pulleys), unless specified by physician to progress more slowly
- 2. Joint mobilization for scapular and glenohumeral mobility
- 3. Gentle exercises for scapular stabilizers and rotator cuff May include sub-max isometrics, PNF, and supervised UBE
- 4. Progress AROM with education for quality of motion to avoid shoulder hiking

*Patient must be able to elevate without hiking shoulder to progress

12-18 Weeks RESISTIVE EXERCISES HELD AS PER SURGEON INSTRUCTIONS